

I. LDA TOTAL ENERGY

Total energy as given by Singh [1] eq. (2.1.3) is

$$E_0 = T_{S,0} + E_{ei} + E_H + E_{ii} + E_{XC} \quad (1)$$

where:

$T_{S,0}$ is the kinetic energy,

E_{ei} is Coulomb electron-nuclei interaction,

E_H is the Hartree energy

$$E_H = \frac{e^2}{2} \int d\vec{r} d\vec{r}' \frac{\rho(\vec{r})\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad (2)$$

E_{ii} is the nuclei-nuclei Coulomb interaction, and
 E_{XC} is the exchange correlation energy

Corresponding Kohn-Sham equation is (Singh eqs. 2.1.6-2.1.9)

$$(T + V_{ei} + V_H + V_{XC})\varphi_i = \epsilon_i\varphi_i \quad (3)$$

with

$$V_H(\vec{r}) = e^2 \int d\vec{r}' \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad (4)$$

$$V_{XC}(\vec{r}) = \frac{\delta E_{XC}}{\delta \rho} \quad (5)$$

and

$$\rho = \sum_{occ} \varphi_i^* \varphi_i \quad (6)$$

To express the kinetic energy $T_{S,0}$ in (1) using $\sum_{occ} \epsilon_i$, equation (3) is multiplied from left by φ_i^* , integrated over \vec{r} and summed over occupied orbitals. The result is:

$$T_{S,0} = \sum_{occ} \epsilon_i - \int d\vec{r} \rho(\vec{r}) [V_{ei} + V_H + V_{XC}] \quad (7)$$

This expression is then substituted to (1). Taking into account that

$$E_{ei} = \int d\vec{r} \rho(\vec{r}) V_{ei}; \quad E_H = \frac{1}{2} \int d\vec{r} \rho(\vec{r}) V_H \quad (8)$$

$$\tilde{E}_0 = \sum_{occ} \epsilon_i + E_{ii} + E_{XC} - \int d\vec{r} \rho(\vec{r}) \left[\frac{1}{2} V_H + V_{XC} \right] \quad (9)$$

which is the desired expression (Singh 2.2.2). The tilde upon E_0 means that $T_{S,0}$ was replaced by $\sum_{occ} \epsilon_i$. The above derivation assumes the non-spin polarized calculation, but the generalization for the spin-polarized problems is straightforward and obvious.

II. LDA+U TOTAL ENERGY

First we assume (general case will be treated afterwards) that:

- Density matrix is diagonal.
- U is the same for all Coulomb interaction ($U_{ij} \equiv U$) and J is the same for all exchange interaction ($J_{ij} \equiv J$).

In the LDA+U methods the total energy may be written as:

$$E = E_0 + E_{LDA+U} \quad (10)$$

A. LDA+U^{DFT}

In the AMF (Around Mean Field) method of Czyzyk and Sawatzky [2] which is also the method we prefer now and call LDA+U^{DFT}:

$$E_{LDA+U} = -\frac{U-J}{2} \sum_{iat} \sum_{m=-l}^l \sum_{\sigma=\uparrow,\downarrow} (n_{m,\sigma} - \bar{n}_\sigma)^2 \quad (11)$$

where the first sum is over atoms selected for the LDA+U correction, $n_{m,\sigma}$ is occupation of the $|l, m, \sigma\rangle$ orbital, and

$$\bar{n}_\sigma = \frac{1}{2l+1} \sum_{m=-l}^l n_{m,\sigma} \quad (12)$$

is the average number of electrons in orbital with spin σ .

The potential which corresponds to E_{LDA+U} may be written as

$$V_{LDA+U} = \sum_{iat} \sum_{m,\sigma} |m,\sigma\rangle v_{m,\sigma} \langle m,\sigma| \quad (13)$$

with

$$v_{m,\sigma} = \frac{\partial E_{LDA+U}}{\partial n_{m,\sigma}} = -(U-J)(n_{m,\sigma} - \bar{n}_\sigma). \quad (14)$$

Kohn-Sham equation is now:

$$(T + V_{ei} + V_H + V_{XC} + V_{LDA+U})\varphi_i = \epsilon_i\varphi_i. \quad (15)$$

The kinetic energy in presence of the LDA+U potential is:

$$T_S = T_{S,0} - \sum_{occ} \langle \varphi_i | V_{LDA+U} | \varphi_i \rangle. \quad (16)$$

Taking into account that V_{LDA+U} is nonzero only within the atomic spheres of selected atoms:

$$T_S = T_{S,0} - \sum_{m,\sigma} \sum_{occ} |\langle \varphi_i | m, \sigma \rangle|^2 v_{m,\sigma} = T_{S,0} - \sum_{m,\sigma} n_{m,\sigma} v_{m,\sigma} \quad (17)$$

The expression for the total energy is now:

$$\tilde{E} = \tilde{E}_0 + E_{LDA+U} - \sum_{m,\sigma} n_{m,\sigma} v_{m,\sigma} \quad (18)$$

Finally, inserting (11),(14) in (18) we get

$$\tilde{E} = \tilde{E}_0 + \frac{U-J}{2} \sum_{m,\sigma} (n_{m,\sigma}^2 - \bar{n}_\sigma^2) \quad (19)$$

which is the same as:

$$\tilde{E} = \tilde{E}_0 + \frac{U-J}{2} \sum_{m,\sigma} (n_{m,\sigma} - \bar{n}_\sigma)^2 \quad (20)$$

Note that the results is an analog of the double correction term for the Hartree energy (compare eqs. 3, 8). This is not surprising, as it is a consequence of the fact that Hartree energy is quadratic function of density and E_{LDA+U} is quadratic in occupation numbers.

B. LDA+U^{SIC}

In the method we call LDA+U^{SIC} (Anisimov et al. [3], Solovyev et al. [4], Liechtenstein et al. [5], Shick et al. [6], also 'atomic limit' version of Czyzyk and Sawatzky [2]):

$$E_{LDA+U} = E^{ee} - E^{dc} \quad (21)$$

where E_{ee} is the mean field approximation to the electron-electron interaction in the spherically symmetrical atom, E^{dc} is the double counting correction. Using the same approximations as in preceding subsection:

$$E^{ee} = \frac{U}{2} N^2 - \frac{J}{2} \sum_{\sigma} N_{\sigma}^2 - \frac{U-J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 \quad (22)$$

where N is the total number of electrons, N_{σ} is the total number of electrons with spin σ :

$$N_{\sigma} = \sum_m n_{m,\sigma}; \quad N = N_{\uparrow} + N_{\downarrow}. \quad (23)$$

The double summation term is:

$$E^{dc} = \frac{U}{2} N(N-1) - \frac{J}{2} \sum_{\sigma} N_{\sigma}(N_{\sigma}-1) \quad (24)$$

and

$$E_{LDA+U} = \frac{U-J}{2} (N - \sum_{m,\sigma} n_{m,\sigma}^2). \quad (25)$$

additional potential, which corresponds to this energy is given by (13) with:

$$v_{m,\sigma} = \frac{U-J}{2} (1 - 2n_{m,\sigma}) \quad (26)$$

Proceeding as in the above subsection we get:

$$\tilde{E} = \tilde{E}_0 + \frac{U-J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 \quad (27)$$

If $U_{ij} \equiv U$, $J_{ij} \equiv J$ and $n_{i,j,\sigma} \equiv n_{i,\sigma} \delta_{i,j}$ is substituted in eq. 23 of Shick et al. [6], identical result is obtained.

C. LDA+U^{DFT} - general case

To simplify the equations we introduce the notation:

$$\tilde{n}_{m,m',\sigma} = n_{m,m',\sigma} - \bar{n}_{m,m,\sigma} \delta_{m,m'} \quad (28)$$

where $n_{m,m',\sigma}$ is (m, m') element of the occupation number matrix with spin σ . E_{LDA+U} is then

$$E_{LDA+U} = \frac{1}{2} \sum_{m_1, m_2, m_3, m_4}^{\sigma, \sigma'} \tilde{n}_{m_1, m_2, \sigma} \tilde{n}_{m_3, m_4, \sigma'} \{ \langle m_1, m_3 | V_{ee} | m_2, m_4 \rangle - \langle m_1, m_3 | V_{ee} | m_4, m_2 \rangle \delta_{\sigma, \sigma'} \} \quad (29)$$

with

$$\begin{aligned} \langle m_1, m_3 | V_{ee} | m_2, m_4 \rangle &= \sum_k a_k(m_1, m_2, m_3, m_4) F_k; \\ a_k(m_1, m_2, m_3, m_4) &= \\ &= \frac{4\pi}{2k+1} \sum_{q=-k}^k \langle l m_1 | Y_{kq} | l m_2 \rangle \langle l m_3 | Y_{kq}^* | l m_4 \rangle \end{aligned} \quad (30)$$

The potential (14) now becomes a matrix with its (m, m') element:

$$v_{m_1, m_2, \sigma} = \sum_{m_3, m_4, \sigma'} \tilde{n}_{m_3, m_4, \sigma'}$$

$$\{ \langle m_1, m_3 | V_{ee} | m_2, m_4 \rangle - \langle m_1, m_3 | V_{ee} | m_4, m_2 \rangle \delta_{\sigma, \sigma'} \} \quad (31)$$

The expression for the kinetic energy is:

$$T_S = T_{S,0} - \sum_{m_1, m_2, \sigma} v_{m_1, m_2, \sigma} \sum_{occ} \langle \varphi_i | m_1, \sigma \rangle \langle m_2, \sigma | \varphi_i \rangle \quad (32)$$

Noting that

$$\sum_{occ} \langle \varphi_i | m_1, \sigma \rangle \langle m_2, \sigma | \varphi_i \rangle = n_{m_2, m_1, \sigma} \quad (33)$$

the total energy, in which kinetic energy was replaced by the sum over eigenvalues, is rewritten as:

$$\tilde{E} = \tilde{E}_0 + E_{LDA+U} - \sum_{\sigma} Tr \{ n_{\sigma} v_{\sigma} \} \quad (34)$$

which is the final result to be used to calculate the total energy in the rotationally invariant LDA+U^{DFT} scheme.

Expression (34) should be correct whatever is the form of the potential v_σ . As the trace is invariant with respect to the unitary transformation, the representation in which $n_{m,m',\sigma}$ is diagonal may be used, without loss of generality. Otherwise we follow Shick et al. [6] (SLP in what follows), as only in this paper the expression for \tilde{E} is given. To make connection with above sections we denote:

$$\begin{aligned} U_{mm'} &= U + u_{mm'}; \\ J_{mm'} &= J + j_{mm'}, m \neq j, \\ J_{mm'} &= U + j_{mm'}, m = j, j_{mm} = u_{mm} \end{aligned} \quad (35)$$

The energy E^{ee} (eq. 22, SLP eq. 2) is then

$$E^{ee} = \frac{1}{2} \sum_{m,m'}^{\sigma,\sigma'} \{U_{mm'} - J_{mm'} \delta_{\sigma,\sigma'}\} n_{m,\sigma} n_{m',\sigma'} \quad (36)$$

Using (35) this becomes

$$E^{ee} = \frac{U}{2} N^2 - \frac{J}{2} \sum_{\sigma} N_{\sigma}^2 - \frac{U-J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 + e \quad (37)$$

where e is given by ($e=0$, if $u_{m,m'} = j_{m,m'} = 0$)

$$e = \frac{1}{2} \sum_{m,m'}^{\sigma,\sigma'} \{u_{mm'} - j_{mm'} \delta_{\sigma,\sigma'}\} n_{m,\sigma} n_{m',\sigma'} \quad (38)$$

Double counting energy is still given by (24) so that

$$E^{ee} - E^{dc} = \frac{N}{2} (U - J) - \frac{U - J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 + e \quad (39)$$

According to eqs. (23-24) of SLP the energy \tilde{E} is:

$$\tilde{E} = \tilde{E}_0 - E^{ee} + E^{dc} + \frac{1}{2} (U - J) N \quad (40)$$

inserting (39) we get

$$\tilde{E} = \tilde{E}_0 + \frac{U - J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 - e \quad (41)$$

Now we come to our derivation. The potential (SLP eq. 16) is:

$$\begin{aligned} v_{m,\sigma} &= \sum_{m',\sigma'} (U_{m,m'} - J_{m,m'} \delta_{\sigma,\sigma'}) n_{m',\sigma'} \\ &\quad - U(N - \frac{1}{2}) + J(N_{\sigma} - \frac{1}{2}) \end{aligned} \quad (42)$$

Using the same notation as above

$$v_{m,\sigma} = (U - J) (\frac{1}{2} - n_{m,\sigma}) + w_{m,\sigma} \quad (43)$$

where

$$w_{m,\sigma} = \sum_{m',\sigma'} (u_{m,m'} - j_{m,m'} \delta_{\sigma,\sigma'}) n_{m',\sigma'} \quad (44)$$

trace of $(v_\sigma n_\sigma)$ is

$$\text{Tr}(v_\sigma n_\sigma) = (U - J) (\frac{1}{2} N_\sigma - \sum_m n_{m,\sigma}^2) + \text{Tr}(w_\sigma n_\sigma) \quad (45)$$

So that using (34) we get the result

$$\begin{aligned} \tilde{E} &= \tilde{E}_0 + \frac{N}{2} (U - J) \\ &\quad - \frac{U - J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 + e \\ &\quad - \frac{N}{2} (U - J) + (U - J) \sum_{m,\sigma} n_{m,\sigma}^2 \\ &\quad - \sum_{\sigma} \text{Tr}(w_\sigma n_\sigma) \end{aligned} \quad (46)$$

which reduces to

$$\tilde{E} = \tilde{E}_0 + \frac{U - J}{2} \sum_{m,\sigma} n_{m,\sigma}^2 + e - \sum_{\sigma} \text{Tr}(w_\sigma n_\sigma) \quad (47)$$

Noting that

$$e = 2 \sum_{\sigma} \text{Tr}(w_\sigma n_\sigma) \quad (48)$$

we get the same result as SLP (see eq. 41, SLP eq. 23-24).

III. CONCLUSION

To calculate the total energy in the LDA+U method, in case the sum over eigenvalues is used instead of the kinetic energy, we recommend to use the equation:

$$\tilde{E} = \tilde{E}_0 + E_{LDA+U} - \sum_{\sigma} \text{Tr}\{n_\sigma v_\sigma\} \quad (49)$$

This formula holds for any LDA+U potential which may be written in the form

$$V_{LDA+U} = \sum_{iat} \sum_{m,m',\sigma} |m,\sigma\rangle v_{m,m',\sigma} \langle m',\sigma|, \quad (50)$$

in particular for both 'DFT' and 'SIC' methods. Moreover, it is rotationally invariant and may be easily implemented.

- [1] D.J. Singh, *Planewaves, pseudopotentials and the LAPW method* (Kluwer Academic, Dordrecht, 1994).
- [2] M.T. Czyzyk and G.A. Sawatzky, Phys. Rev. B **49**, 14211 (1994).
- [3] V.I. Anisimov, I.V. Solovyev, M.A. Korotin, M.T. Czyzyk, and G.A. Sawatzky, Phys. Rev. B **48**, 16929 (1993).
- [4] I.V. Solovyev, P.H. Dederichs, and V.I. Anisimov, Phys. Rev. B **50**, 16861 (1994).
- [5] A.I. Liechtenstein, V.I. Anisimov, and J. Zaanen, Phys. Rev. B **52**, 5467 (1995).
- [6] A.B. Shick, A.I. Liechtenstein, and W.E. Pickett, Phys. Rev. B **60**, 10763 (1999).